arXiv:1211.7358v1 [cond-mat.stat-mech] 30 Nov 2012

Correlation functions in one-dimensional spin lattices with Ising and Heisenberg bonds

Stefano Bellucci[1] and Vadim Ohanyan[2,3]
[1]*INFN-Laboratori Nazionali di Frascati, Via E. Fermi 40, 00044 Frascati, Italy*
[2]*Department of Theoretical Physics, Yerevan State University, Alex Manoogian 1, 0025 Yerevan, Armenia*
[3]*ICTP, Strada Costiera 11, I-34151 Trieste, Italy*

(Dated: December 3, 2012)

A general technique of exact calculation of any correlation functions for the special class of one-dimensional spin models containing small clusters of quantum spins assembled to a chain by alternating with the single Ising spins is proposed. The technique is a natural generalization of that in the models solved by a classical transfer matrices. The general expressions for corresponding matrix operators which are the key components of the technique are obtained. Despite the spatial inhomogeneity, such class of spin chains is proven to have only one correlation length for all types of correlation functions. The technique is illustrated on two examples: symmetric diamond chain and asymmetric sawtooth chain.

PACS numbers: 75.10.Jm
Keywords: Heisenberg-Ising chains, correlation functions, classical transfer matrix, diamond chain, sawtooth chain

I. INTRODUCTION

Recently much attention has been paid to the exactly solvable models of one-dimensional magnetism, consisting of the small quantum spin clusters connected to each other via Ising spins.[1–26] For the sake of simplicity we will call them Heisenberg–Ising chains (HIC). The interest toward such kind of spin systems stems out from the several features they possess. First of all, they allow one to obtain an exact statistical-mechanical solution of the problem of strongly interacting spins in terms of classical transfer matrix[27–30]. In addition to that, such systems combine the quantum and classical properties and also, to some extent, are connected to the problems and models of molecular magnetism[31], as the classical transfer matrix is constructed by the diagonalization of the small quantum spin cluster. And finally, very recently the family of trimetallic coordination polymer compounds has been sythesized[32–34]. These compounds in the magnetic sense are indeed a one-dimensional magnets with both classical (Ising) and quantum (Heisenberg) bonds. The single-chain magnet reported in Ref. 32 is an example of a Heisenberg-Ising chain of triangles with three-spin linear quantum clusters; another interesting molecular magnet (spin ring) based on the highly anisotropic properties of the Dy^{3+} ions has been reported in Ref. 33. Both systems have been analyzed by the transfer-matrix solution of Heisenberg-Ising spin chains in Ref. 34. Another class of examples is given by chain magnets with two alternating magnetic ions, thus giving rise to alternating quantum and classical spins (see, e.g., Refs. 35–37). Let us describe the general scheme of the HIC:

- There is a small cluster of a few quantum spins with a certain topology of bonds which could contain various types of spin-spin interaction as well as interactions with external fields. The Hamiltonian of the single small cluster is, thus, a finite dimensional matrix.

- These clusters are assembled to the chain or any other one-dimensional structure by alternating with the single Ising spins.

- The interaction between the single Ising spins and any quantum spin from small cluster must contain only z-components (Ising interaction).

For the mixed quantum–classical spin system with the above mentioned properties the exact solution is obtained in the following way. The idea is to apply a classical transfer-matrix formalism in which the entries of classical transfer matrix is obtained from the diagonalization of a small quantum spin cluster. Although the number of paper devoted to the exact solutions of the HIC and to their magneto-thermal properties is rather impressive, only few models have been solved within the classical transfer matrix formalism[5,9–11,13–17,19–22,26], while the other method of so-called decoration iteration transformation has been also used[1–4,8,12,24,25]. Particularly, within the direct classical transfer matrix technique the following models have been solved exactly: the double tetrahedra chain with quantum triangles of spins[9,11], the sawtooth chain with asymmetric pair of quantum spins[10,13], diamond chain[14,16,17,20–22,26] and orthogonal-dimer chain with anisotropic and asymmetric quantum triangles[19]. Despite the complete and exact description of the ground states and ground states phase diagrams, all thermodynamic functions, finite-temperature entropy and magnetization, magnetocaloric effect, etc, the expressions for the correlation function are still missing. In the present paper we are going to fill this gap by developing the exact technique for the calculation of any finite-temperature correlation function for any HIC.

The paper is organized as follows. In the second Section we remind the reader the main features of the classical transfer-matrix formalism for the HIC, in the next section we present the formalism of calculating the correlation function of HIC in terms of traces of certain matrix product and proof that for all kinds of correlation functions the correlation length is the same. In the Section IV we apply the general formalism for several models. The paper ends with a Conclusion.

II. CLASSICAL TRANSFER-MATRIX FORMALISM FOR HIC

The Hamiltonian of a HIC has the form of a sum of a local block-Hamiltonians $\mathcal{H}_n$ which correspond to the n-th small quantum cluster and its interaction with left and right Ising spins:

$$\mathcal{H} = \sum_{n=1}^{N} \left(\mathcal{H}_n - \frac{H}{2} \left(\sigma_n + \sigma_{n+1} \right) \right), \tag{2.1}$$

here σ_n is the Ising spin situated between n-th and (n+1)-th small cluster and H is the magnetic field acting on Ising spins. Due to the properties listed above the block Hamiltonians commute to each other,

$$[\mathcal{H}_i, \mathcal{H}_j] = 0. \tag{2.2}$$

This fact allows one to expand the exponential in the expression for the partition function of the model and obtain an expression which by its structure corresponds to the classical transfer matrix scheme:

$$\begin{aligned} \mathcal{Z} &= \mathrm{Tr}_{\mathrm{S}} \sum_{(\sigma)} \sum_{(\tau)} \exp\left(-\beta \mathcal{H}\right) = \sum_{(\sigma)} \prod_{n=1}^{N} e^{-\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \sum_{(\tau)} \mathrm{Tr}_n e^{-\beta \mathcal{H}_n} = \\ & \sum_{(\sigma)} \prod_{n=1}^{N} e^{-\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \sum_{(\tau)} \sum_{i=1}^{L} e^{-\beta \lambda_i (\sigma_n, \sigma_{n+1} | \tau_{n,b})} = \sum_{(\sigma)} \prod_{n=1}^{N} T_{\sigma_n, \sigma_{n+1}} = \mathrm{Tr} \mathbf{T}^N, \end{aligned} \tag{2.3}$$

here the cyclic boundary conditions are supposed, $\lambda_i(\sigma_n, \sigma_{n+1}|\tau_{n,b})$ are the eigenvalues of the block Hamiltonian $\mathcal{H}_n$(which is supposed to be an L by L matrix) depending on the values of classical spins τ and classical spins σ_n which alternate with the blocks, Tr_n means trace over the states of the n-th quantum block, and Tr stands for the trace for the 2 by 2 transfer matrix with entries $T_{\sigma_n,\sigma_{n+1}}$. After summation over the values of τ for each block one gets the expression depending only on σ_n and σ_{n+1} which is the classical 2 by 2 transfer matrix

$$\mathbf{T} = \begin{pmatrix} e^{\beta H} Z_{++} & Z_{+-} \\ Z_{-+} & e^{-\beta H} Z_{--} \end{pmatrix}, \tag{2.4}$$

where $Z_{\sigma_n,\sigma_{n+1}} = \sum_{(\tau)} \sum_{i=1}^{L} e^{-\beta \lambda_i (\sigma_n, \sigma_{n+1} | \tau_{n,b})}$ is the partition function for one block, depending on the values of two neighbor Ising spins, signs "+" and "−" in indexes correspond to $\sigma = 1/2$ and $\sigma = -1/2$ respectively. The formalism of classical transfer-matrix allows one to obtain exact analytic expressions for almost any expectation values of the function of local variables (spins) and, of course, for any thermodynamic functions, as the expression for the free energy per block can be obtained in a straightforward way by finding the maximal eigenvalue of the transfer matrix (2.4):

$$f = -\frac{1}{\beta} \log \frac{1}{2} \left(e^{\beta H} Z_{++} + e^{-\beta H} Z_{--} + \sqrt{\left(e^{\beta H} Z_{++} - e^{-\beta H} Z_{--} \right)^2 + 4 Z_{+-} Z_{-+}} \right). \tag{2.5}$$

From this expression any thermodynamic function, like magnetization, entropy, specific head e.t.c. can be obtained exactly. However, to obtain correlation functions of the spins demands more efforts and further development of the classical transfer-matrix technique.

III. CORRELATION FUNCTIONS AND CORRELATION LENGTH

The block Hamiltonian can contain various terms with quantum and classical spin variables and which are determined by the topology of the elementary block and the character of interactions. Let us denote the quantum spins by $\mathbf{S}$; then one can suppose that block Hamiltonian depends on $\mathbf{S}$, σ and may be other classical spins τ_b,

$$\mathcal{H}_n = \mathcal{H}\left(\mathbf{S}_{n,a}, \sigma_n, \sigma_{n+1}, \tau_{n,b}\right). \tag{3.1}$$

Let us concentrate on the scheme of calculation of the expectation values of combinations of spins operators. First of all, let us start form the simplest case, the expectation values of σ-s, which is quite similar to that of the Ising chain[27]. Thus, for the thermodynamic average of just one σ one obtains

$$\begin{aligned}\langle\sigma_j\rangle &= \frac{1}{\mathcal{Z}}\sum_{(\sigma)}\sigma_j\prod_{n=1}^{N}T_{\sigma_n,\sigma_{n+1}} \\ &= \frac{1}{\mathcal{Z}}\sum_{(\sigma)}T_{\sigma_1,\sigma_2}...\sigma_j T_{\sigma_j,\sigma_{j+1}}...T_{\sigma_N,\sigma_1}. \end{aligned}\tag{3.2}$$

This expression is the trace of the product of N transfer-matrices with the matrix

$$\sigma^z = \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix} \tag{3.3}$$

inserted before the j-th transfer matrix in the trace formula. Thus, we have

$$\langle\sigma_j\rangle = \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\sigma^z\mathbf{T}^N\right) = \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\sigma^z\mathbf{A}\mathbf{\Lambda}^N\mathbf{A}^{-1}\right), \tag{3.4}$$

where we used the matrix identity $\mathbf{T} = \mathbf{A}\mathbf{\Lambda}\mathbf{A}^{-1}$, with the diagonal matrix $\mathbf{\Lambda}$ and the orthogonal matrix $\mathbf{A}$. Thus,

$$\mathbf{\Lambda} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \qquad \mathbf{A} = \begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix}, \tag{3.5}$$

where $\lambda_{1,2}$ are the eigenvalues of the transfer-matrix, and ϕ is a parameter. Having all these notations we now can express the statistical mechanical average from Eq. (3.2) in terms of the entries of the orthogonal matrix diagonalizing the transfer-matrix. Using the properties of the trace (cyclic permutation inside the trace does not change it) one can get

$$\begin{aligned}\langle\sigma_j\rangle &= \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{A}^{-1}\sigma^z\mathbf{A}\mathbf{\Lambda}^N\right) = \\ & \frac{1}{2}\frac{(\cos 2\phi)\lambda_1^2 - (\sin 2\phi)\lambda_2^N}{\lambda_1^N + \lambda_2^N}\end{aligned} \tag{3.6}$$

Now, we should take care about the thermodynamic limit ($N \to \infty$). It is easy to see, that in this limit Eq. (3.6) leads to (suppose that λ_1 is greater than λ_2)

$$\langle\sigma_j\rangle = \frac{1}{2}\cos 2\phi \equiv \frac{1}{2}M_\sigma, \tag{3.7}$$

where M_σ is the partial magnetization corresponding to the subsystem with σ-spins,

$$M_\sigma = \frac{e^{\beta H}Z_{++} - e^{-\beta H}Z_{--}}{\sqrt{\left(e^{\beta H}Z_{++} - e^{-\beta H}Z_{--}\right)^2 + 4Z_{+-}Z_{-+}}} \tag{3.8}$$

Thus, we can express the entries of the orthogonal matrix diagonalising the transfer matrix in terms of the subsystem magnetization M_σ:

$$\mathbf{A} = \frac{1}{\sqrt{2}}\begin{pmatrix} \sqrt{1+M_\sigma} & -\sqrt{1-M_\sigma} \\ \sqrt{1-M_\sigma} & \sqrt{1+M_\sigma} \end{pmatrix} \tag{3.9}$$

These quantities are the building blocks for calculation of all correlation function for the HIC. As will be clear from the further consideration, any correlation function of HIC has the form of the trace of a certain product of transfer matrices and special matrices representing the spin operators. Moreover, these matrices are constructed from the local expectation values of spin operators with the aid of the matrix $\mathbf{A}$ in the standard way $\widetilde{\mathbf{P}} \equiv \mathbf{A}^{-1}\mathbf{P}\mathbf{A}$. Let us write down the entries of a general matrix of that type

$$\begin{aligned}
\widetilde{P}_{++} &= \frac{1}{2}\left[(1+M_\sigma)P_{++} + \sqrt{1-M_\sigma^2}\left(P_{+-}+P_{-+}\right) + (1-M_\sigma)P_{--}\right], \\
\widetilde{P}_{+-} &= \frac{1}{2}\left[\sqrt{1-M_\sigma^2}\left(P_{--}-P_{++}\right) + (1+M_\sigma)P_{+-} - (1-M_\sigma)P_{-+}\right], \\
\widetilde{P}_{-+} &= \frac{1}{2}\left[\sqrt{1-M_\sigma^2}\left(P_{--}-P_{++}\right) + (1+M_\sigma)P_{-+} - (1-M_\sigma)P_{+-}\right], \\
\widetilde{P}_{--} &= \frac{1}{2}\left[(1-M_\sigma)P_{++} - \sqrt{1-M_\sigma^2}\left(P_{+-}+P_{-+}\right) + (1+M_\sigma)P_{--}\right].
\end{aligned} \tag{3.10}$$

For instance, the matrix $\widetilde{\sigma}^z = \mathbf{A}^{-1}\sigma_z\mathbf{A}$ which is the building block for constructing various correlation functions of σ_j has a particular simple form:

$$\widetilde{\sigma}^z = \frac{1}{2}\begin{pmatrix} M_\sigma & -\sqrt{1-M_\sigma^2} \\ -\sqrt{1-M_\sigma^2} & -M_\sigma \end{pmatrix} \tag{3.11}$$

The simple two-site correlation function is just a thermal average of the product of two σ spins situated at a distance r form each other:

$$\langle \sigma_j \sigma_{j+r} \rangle = \frac{1}{\mathcal{Z}} \sum_{(\sigma)} \sigma_j \sigma_{j+r} \prod_{n=1}^{N} T_{\sigma_n, \sigma_{n+1}} = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r} \sigma^z \mathbf{T}^r \sigma^z \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r} \widetilde{\sigma}^z \mathbf{\Lambda}^r \widetilde{\sigma}^z \right).$$

Thus, using the matrix form of $\widetilde{\sigma}^z$ and taking a thermodynamic limit one obtains:

$$\begin{aligned} \langle \sigma_j \sigma_{j+r} \rangle &= \frac{1}{4} \left(M_\sigma^2 + \left(1 - M_\sigma^2\right) e^{-\frac{r}{\xi}} \right), \qquad (3.12) \\ \xi &= \frac{1}{\log\left(\frac{\lambda_1}{\lambda_2}\right)}, \end{aligned}$$

where ξ is a correlation length. So far, all results have the same form as those for a simple Ising chain, the difference is only in the form of the magnetization for σ-spins, which in our case is given by Eqs. (3.8) and (2.4). Now, let us turn to the more complicated correlation functions of the quantum spins with each other and mixed correlation functions,

$$\begin{aligned} \langle S_{j,a}^{\alpha} S_{j+r,b}^{\beta} \rangle &= \frac{1}{\mathcal{Z}} \mathbf{Tr}_S \sum_{(\sigma)} \sum_{(\tau)} S_{j,a}^{\alpha} S_{j+r,b}^{\beta} \prod_{n=1}^{N} e^{-\beta \mathcal{H}_n} e^{\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \qquad (3.13) \\ &= \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r-1} \mathbf{\Sigma}_a^{\alpha} \mathbf{T}^{r-1} \mathbf{\Sigma}_b^{\beta} \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r-1} \widetilde{\mathbf{\Sigma}}_a^{\alpha} \mathbf{\Lambda}^{r-1} \widetilde{\mathbf{\Sigma}}_b^{\beta} \right), \\ \langle S_{j,a}^{\alpha} \sigma_{j+r} \rangle &= \frac{1}{\mathcal{Z}} \mathbf{Tr}_S \sum_{(\sigma)} \sum_{(\tau)} S_{j,a}^{\alpha} \sigma_{j+r} \prod_{n=1}^{N} e^{-\beta \mathcal{H}_n} e^{\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \\ &= \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r-1} \mathbf{\Sigma}_a^{\alpha} \mathbf{T}^{r} \sigma^z \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r-1} \widetilde{\mathbf{\Sigma}}_a^{\alpha} \mathbf{\Lambda}^{r} \widetilde{\sigma}^z \right) \\ \langle S_{j,a}^{\alpha} \tau_{j+r,b} \rangle &= \frac{1}{\mathcal{Z}} \mathbf{Tr}_S \sum_{(\sigma)} \sum_{(\tau)} S_{j,a}^{\alpha} \tau_{j+r,b} \prod_{n=1}^{N} e^{-\beta \mathcal{H}_n} e^{\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \\ &= \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r-1} \mathbf{\Sigma}_a^{\alpha} \mathbf{T}^{r-1} \tau_b \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r-1} \widetilde{\mathbf{\Sigma}}_a^{\alpha} \mathbf{\Lambda}^{r-1} \widetilde{\tau}_b \right), \\ \langle \tau_{j,a} \tau_{j+r,b} \rangle &= \frac{1}{\mathcal{Z}} \mathbf{Tr}_S \sum_{(\sigma)} \sum_{(\tau)} \tau_{j,a} \tau_{j+r,b} \prod_{n=1}^{N} e^{-\beta \mathcal{H}_n} e^{\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \\ &= \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r-1} \tau_a \mathbf{T}^{r-1} \tau_b \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r-1} \widetilde{\tau}_a \mathbf{\Lambda}^{r-1} \widetilde{\tau}_b \right), \\ \langle \tau_{j,a} \sigma_{j+r} \rangle &= \frac{1}{\mathcal{Z}} \mathbf{Tr}_S \sum_{(\sigma)} \sum_{(\tau)} \tau_{j,a} \sigma_{j+r} \prod_{n=1}^{N} e^{-\beta \mathcal{H}_n} e^{\beta \frac{H}{2} (\sigma_n + \sigma_{n+1})} \\ &= \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{T}^{N-r-1} \tau_a \mathbf{T}^{r} \sigma \right) = \frac{1}{\mathcal{Z}} \mathbf{Tr} \left(\mathbf{\Lambda}^{N-r-1} \widetilde{\tau}_a \mathbf{\Lambda}^{r} \widetilde{\sigma} \right), \end{aligned}$$

where the new 2 by 2 matrices $\mathbf{\Sigma}_a^{\alpha}$ and τ_a are the classical transfer-matrix formalism representation of the α-component of the a-th quantum spin and classical a-th spins from the corresponding quantum clusters. Their entries are found according to

$$\begin{aligned}\left(\Sigma_a^\alpha\right)_{\sigma_j,\sigma_{j+1}} &= e^{\beta\frac{H}{2}(\sigma_j+\sigma_{j+1})}\sum_{(\tau_j)}\mathbf{Tr}_j\left(S^\alpha_{j,a}e^{-\beta\mathcal{H}_j}\right)\\ \left(\tau_a\right)_{\sigma_j,\sigma_{j+1}} &= e^{\beta\frac{H}{2}(\sigma_j+\sigma_{j+1})}\sum_{(\tau_j)}\tau_{j,a}\mathbf{Tr}_j e^{-\beta\mathcal{H}_j} \quad (3.14)\end{aligned}$$

As the expressions under the trace in Eq.(3.13) have the same structure, let us write down a general expression for such a correlation function

$$\begin{aligned}&\frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{\Lambda}^{N-r-1}\widetilde{\mathbf{P}}\mathbf{\Lambda}^{r-1}\widetilde{\mathbf{R}}\right) \quad (3.15)\\ &= \frac{1}{\lambda_1^2}\widetilde{P}_{++}\widetilde{R}_{++} + \frac{1}{\lambda_1\lambda_2}\widetilde{P}_{+-}\widetilde{R}_{-+}e^{-r/\xi},\end{aligned}$$

where the correlation length ξ is the same as in Eq.(3.12). Obviously, the part independent of r is the product of expectation values of quantities corresponding to matrices $\mathbf{P}$ and $\mathbf{R}$,

$$\langle P_j\rangle = \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{\Lambda}^{N-1}\widetilde{\mathbf{P}}\right) = \frac{1}{\lambda_1}\widetilde{P}_{++}. \quad (3.16)$$

The product of two eigenvalues of the 2 by 2 transfer matrix is its determinant, thus, eventually one can write down the following general form of pair-correlation function for HIC:

$$\begin{aligned}\langle P_j R_{j+r}\rangle &= \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{\Lambda}^{N-r-1}\widetilde{\mathbf{P}}\mathbf{\Lambda}^{r-1}\widetilde{\mathbf{R}}\right) \quad (3.17)\\ &= \langle P_j\rangle\langle R_j\rangle + \frac{\widetilde{P}_{+-}\widetilde{R}_{-+}}{\det\mathbf{T}}e^{-r/\xi}\end{aligned}$$

If one of the matrices is σ the result is different, because the corresponding expression contains one more transfer matrix,

$$\begin{aligned}&\frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{\Lambda}^{N-r-1}\widetilde{\mathbf{P}}\mathbf{\Lambda}^{r}\widetilde{\sigma}\right) \quad (3.18)\\ &= \frac{1}{2\lambda_1}\left(\widetilde{P}_{++}M_\sigma - \widetilde{P}_{+-}\sqrt{1-M_\sigma^2}e^{-r/\xi}\right).\end{aligned}$$

The most important conclusion one can make from this general consideration is the existence of only one correlation length for the any kind of HIC. Despite the inhomogeneous nature of such kind of systems, the decay of correlations between various spins belonging to different blocks is characterized by the same correlation length, defined by the ratio of the maximal and the next to the maximal(in our case of 2 by 2 transfer matrix, just the "first" and the "second") eigenvalues of transfer matrix. This is, however, the true only for a long range correlation with respect to the block structure of the system; inside one block short range correlations are non-uniform.

IV. EXAMPLES

In this section we are going to illustrate the technique of calculation of correlation functions for HIC on several examples. As the technique, despite its clearness and simplicity, leads to rather cumbersome expressions, we will present explicit calculation for the simplest models which have been investigated earlier in a context of magnetic and thermodynamic properties[1–26]. One can distinguish at least two classes of HIC: the models where each block has a left-right symmetry with respect to interaction with σ spins[1–9,11,14], which implies that the term describing the interaction between block spins and their left and right σ-neighbors has the form $K\left(\sigma_j+\sigma_{j+1}\right)\sum_a S^z_{j,a}$, which is completely symmetric with respect to the permutation of the operators $S^z_{j,a}$, and the models with are non symmetric with respect to left and right σ-spins on each block[10,13,19]. It is worth mentioning, that the experimentally obtained single chain magnet compound with Ising and Heisenberg bonds, which is an existing example of HIC, belongs to the second type[32,34]. Magnetic properties of the asymmetric model are vast and complicated, they can exhibit a large number of ground states with a breaking of translational symmetry or doubling of the unit cell(block)[10,13,19]. In the most symmetric case the block Hamiltonian is completely symmetric with respect to permutation of all spins in it. Such models are the simplest ones. For instance, the various variants of the Heisenberg–Ising diamond chain[4,8,14] and Heisenebrg-Ising tetrahedral[5] and double-tetrahedral chains[9,11] belong to this class of HIC. It is obvious, that in this case the matrices $\mathbf{\Sigma}^\alpha_\mathbf{a}$ are the same for all spins inside one block, or, in other words, and in this case one does not need the additional index a.



FIG. 1: Heisenberg-Ising diamond chain. Solid (dashed) lines denote quantum XXZ (classical Ising)interaction bonds. All dashed bonds correspond to coupling constant K, the interaction between quantum spins is characterized by coupling constant J and anisotropy Δ.

A. Symmetric diamond-chain

The symmetric Heisenberg-Ising diamond chain is the simplest example of HIC (See Figure 1). Its block Hamiltonian has the form:

$$\begin{aligned}\mathcal{H}_n^{DC} &= J\left(\Delta\left(S^x_{n,1}S^x_{n,2}+S^y_{n,1}S^y_{n,2}\right)+S^z_{n,1}S^z_{n,2}\right)\\ &-\left(H-K\left(\sigma_n+\sigma_{n+1}\right)\right)\left(S^z_{n,1}+S^z_{n,2}\right) \quad (4.1)\end{aligned}$$

The corresponding transfer-matrix is

$$\mathbf{T}^{DC} = 2e^{-\beta\frac{J}{4}} \begin{pmatrix} e^{\beta\frac{H}{2}}\left(\mathrm{ch}\left[\beta\left(H-K\right)\right] + Q^{DC}\right) & \mathrm{ch}\left(\beta H\right) + Q^{DC} \\ \mathrm{ch}\left(\beta H\right) + Q^{DC} & e^{-\beta\frac{H}{2}}\left(\mathrm{ch}\left[\beta\left(H+K\right)\right] + Q^{DC}\right) \end{pmatrix}, \tag{4.2}$$

where

$$Q^{DC} = e^{\beta\frac{J}{2}}\,\mathrm{ch}\left(\beta\frac{J\Delta}{2}\right). \tag{4.3}$$

As the block Hamiltonian is SO(2)-invariant with respect to the z-axis the corresponding $\mathbf{\Sigma}_a^{\alpha}$-matrix is non-zero only for $\alpha = z$ and, as discussed above, in virtue of the symmetry with respect to permutation of $\mathbf{S}$-operators for one block $\mathbf{\Sigma}_1^z = \mathbf{\Sigma}_2^z \equiv \mathbf{\Sigma}^z$. The matrix $\mathbf{\Sigma}^z$ has a rather simple form:

$$\mathbf{\Sigma}^z = \tag{4.4}$$
$$e^{-\beta\frac{J}{4}} \begin{pmatrix} e^{\beta\frac{H}{2}}\,\mathrm{sh}\left[\beta\left(H-K\right)\right] & \mathrm{sh}\left(\beta H\right) \\ \mathrm{sh}\left(\beta H\right) & e^{-\beta\frac{H}{2}}\,\mathrm{sh}\left[\beta\left(H+K\right)\right] \end{pmatrix}$$

According to Eqs.(3.10)-(3.18), the non-zero two-site correlation functions for this model are

$$\langle S_{j,a}^z S_{j+r,b}^z\rangle - \langle S_{j,a}^z\rangle^2 = \frac{\left(2\,\mathrm{sh}\left(\beta H\right)M_\sigma + \left[e^{\beta K}\,\mathrm{ch}\left(\beta\frac{H}{2}\right) - e^{-\beta K}\,\mathrm{ch}\left(\beta\frac{3H}{2}\right)\right]\sqrt{1-M_\sigma^2}\right)^2}{16\left(\mathrm{sh}^2\left(\beta K\right) + 2Q^{DC}\left(\mathrm{ch}\left(\beta K\right)-1\right)\mathrm{ch}\left(\beta H\right)\right)} e^{-\frac{r}{\xi}}, \tag{4.5}$$
$$\langle S_{j,a}^z \sigma_{j+r}^z\rangle - \langle S_{j,a}^z\rangle\langle\sigma_j\rangle = -\frac{e^{-\beta\frac{J}{4}}\left\{2\,\mathrm{sh}\left(\beta H\right)M_\sigma\sqrt{1-M_\sigma^2} + \left[e^{\beta K}\,\mathrm{ch}\left(\beta\frac{H}{2}\right) - e^{-\beta K}\,\mathrm{ch}\left(\beta\frac{3H}{2}\right)\right]\left(1-M_\sigma^2\right)^2\right\}}{4\lambda_1^{DC}} e^{-\frac{r}{\xi}}$$

where

$$\langle S_{j,a}^z\rangle = \frac{1}{\mathcal{Z}}\mathbf{Tr}\left(\mathbf{\Lambda}^{N-1}\widetilde{\mathbf{\Sigma}^{\mathbf{z}}}\right) = \frac{1}{2}M_S \tag{4.6}$$

is the partial magnetization for the sublattice of quantum spins. M_σ is the magnetization of σ-sublattice obtained according to Eq.3.8 from transfer-matrix (4.2) and λ_1^{DC} is the largest eigenvalue of the transfer-matrix (4.2) given by

$$\begin{aligned}\lambda_1^{DC} = e^{-\beta\frac{J}{4}} \quad & \left\{e^{-\beta K}\,\mathrm{ch}\left(\beta\frac{3H}{2}\right) + \left(2Q^{DC} + e^{\beta K}\right)\mathrm{ch}\left(\beta\frac{H}{2}\right)\right. \\ & \left. + \sqrt{\left[e^{-\beta K}\,\mathrm{sh}\left(\beta\frac{3H}{2}\right) + \left(2Q^{DC} - e^{\beta K}\right)\mathrm{sh}\left(\beta\frac{H}{2}\right)\right]^2 + 4\left(\mathrm{ch}\left(\beta H\right) + Q^{DC}\right)^2}\right\}.\end{aligned} \tag{4.7}$$

B. Asymmetric sawtooth chain

The next example we are going to consider is the Heisenberg-Ising sawtooth chain introduced in Refs. 10 and 13. The lattice is depicted in Figure 2. This model is quite convenient for illustrating the asymmetric role of quantum spins from one block as well as for the demonstration of the role of the additional classical spin included in the block. Here the block Hamiltonian contains two parts, one with quantum operators and another one - without them. Thus, one can write down the block Hamiltonian for the sawtooth HIC with 2 quantum spins in one block in the following way:

$$\begin{aligned}&\mathcal{H}_n^{SC} = \mathcal{H}_n^q + K\sigma_n\tau_n - H\tau_n, \\ &\mathcal{H}_n^q = J\left(\Delta\left(S_{n,1}^x S_{n,2}^x + S_{n,1}^y S_{n,2}^y\right) + S_{n,1}^z S_{n,2}^z\right) \\ &+ \left(H - K\left(\sigma_n + \sigma_{n+1} + \tau_n\right)\right)S_{n,1}^z - \left(H - K\sigma_{n+1}\right)S_{n,2}^z.\end{aligned} \tag{4.8}$$

Here all notations are explained in Figure 2. In contrast to the case considered previously in Refs. 10 and



FIG. 2: Heisenberg-Ising sawtooth chain with a pair of quantum spins in one block. The unit cell consists of 4 spins, $\mathbf{S}_{j,1}, \mathbf{S}_{j,2}, \sigma_j$ and τ_j. All bonds denoted by dashed lines are of Ising type and the corresponding coupling constant is K, while the XXZ-interaction corresponding to quantum bond (solid line) is given by coupling and anisotropy J and Δ.

13, here, for the sake of simplicity, we consider all Ising bonds with the same coupling constant K. As the block Hamiltonian contains also the Ising spin τ, in order to obtain the entries of transfer matrix one needs, after taking the trace over quantum states of the pair of $\mathbf{S}$-spin, to sum over the two possible values of the classical spin τ. The entries of the resulting classical transfer matrix are:

$$\begin{aligned}
T^{SC}_{++} &= 2e^{-\beta\frac{J}{4}}\left\{e^{\beta\left(H-\frac{K}{4}\right)}\left[\mathrm{ch}\left(\beta\left(H-K\right)\right)+Q^{SC}_2\right]+e^{\beta\frac{K}{4}}\left[\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+Q^{SC}_1\right]\right\}, \qquad (4.9)\\
T^{SC}_{+-} &= 2e^{-\beta\frac{J}{4}}\left\{e^{\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta H\right)+Q^{SC}_2\right]+e^{-\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+Q^{SC}_1\right]\right\},\\
T^{SC}_{-+} &= 2e^{-\frac{\beta}{4}}\left\{e^{-\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta H\right)+Q^{SC}_2\right]+e^{\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+Q^{SC}_1\right]\right\},\\
T^{SC}_{--} &= 2e^{-\beta\frac{J}{4}}\left\{e^{-\beta\left(H+\frac{K}{4}\right)}\left[\mathrm{ch}\left(\beta\left(H+K\right)\right)+Q^{SC}_2\right]+e^{\beta\frac{K}{4}}\left[\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+Q^{SC}_1\right]\right\},
\end{aligned}$$

where

$$\begin{aligned}
Q^{SC}_1 &= e^{\beta\frac{J}{2}}\,\mathrm{ch}\left(\beta\frac{J\Delta}{2}\right), \qquad (4.10)\\
Q^{SC}_2 &= e^{\beta\frac{J}{2}}\,\mathrm{ch}\left(\beta\frac{\sqrt{K^2+J^2\Delta^2}}{2}\right).
\end{aligned}$$

The block Hamiltonian (4.8) is not symmetric with respect the to permutation of the quantum spin operators $\mathbf{S}_{n,1}$ and $\mathbf{S}_{n,2}$, thus, there are two matrices representing the z-component of the corresponding spin-operators using for the calculation of the correlation functions. In virtue of the SO(2)-invariance of Eq.(4.8) the corresponding matrices for other spin components identically equal to zero. Let us write down the entries of the $\mathbf{\Sigma}^z_1$ and $\mathbf{\Sigma}^z_2$ matrices for the Heisenberg-Ising sawtooth chain.

$$\begin{aligned}
\left(\mathbf{\Sigma}^z_{1,2}\right)_{++} &= e^{-\beta\frac{J}{4}}\left\{e^{\beta\frac{K}{4}}\,\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+e^{\beta\left(H-\frac{K}{4}\right)}\left(\mathrm{ch}\left(\beta\left(H-K\right)\right)\mp W\right)\right\}, \qquad (4.11)\\
\left(\mathbf{\Sigma}^z_{1,2}\right)_{+-} &= e^{-\beta\frac{J}{4}}\left\{e^{-\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\,\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+e^{\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\left(\mathrm{ch}\left(\beta H\right)\mp W\right)\right\},\\
\left(\mathbf{\Sigma}^z_{1,2}\right)_{-+} &= e^{-\beta\frac{J}{4}}\left\{e^{\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\,\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+e^{-\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\left(\mathrm{ch}\left(\beta H\right)\pm W\right)\right\},\\
\left(\mathbf{\Sigma}^z_{1,2}\right)_{--} &= e^{-\beta\frac{J}{4}}\left\{e^{\beta\frac{K}{4}}\,\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+e^{-\beta\left(H+\frac{K}{4}\right)}\left(\mathrm{ch}\left(\beta\left(H+K\right)\right)\pm W\right)\right\},
\end{aligned}$$

where

$$W = K\frac{e^{\beta\frac{J}{2}}\,\mathrm{sh}\left(\beta\frac{\sqrt{K^2+J^2\Delta^2}}{2}\right)}{\sqrt{K^2+J^2\Delta^2}}. \qquad (4.12)$$

and upper(lower) sings correspond to $\mathbf{\Sigma}^z_1(\mathbf{\Sigma}^z_2)$. The matrix τ in terms of which the expectation values and correlation function of the classical spin from block Hamiltonian are expressed has the following entries:

$$
\begin{aligned}
\tau_{++} &= e^{-\beta\frac{J}{4}}\left\{e^{\beta\left(H-\frac{K}{4}\right)}\left[\mathrm{ch}\left(\beta\left(H-K\right)\right)+Q_2^{SC}\right]-e^{\beta\frac{K}{4}}\left[\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+Q_1^{SC}\right]\right\},\\
\tau_{+-} &= e^{-\beta\frac{J}{4}}\left\{e^{\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta H\right)+Q_2^{SC}\right]-e^{-\frac{\beta}{2}\left(H-\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+Q_1^{SC}\right]\right\},\\
\tau_{-+} &= e^{-\frac{\beta}{4}}\left\{-e^{-\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta H\right)+Q_2^{SC}\right]+e^{\frac{\beta}{2}\left(H+\frac{K}{2}\right)}\left[\mathrm{ch}\left(\beta\left(H-\frac{K}{2}\right)\right)+Q_1^{SC}\right]\right\},\\
\tau_{--} &= e^{-\beta\frac{J}{4}}\left\{-e^{-\beta\left(H+\frac{K}{4}\right)}\left[\mathrm{ch}\left(\beta\left(H+K\right)\right)+Q_2^{SC}\right]+e^{\beta\frac{K}{4}}\left[\mathrm{ch}\left(\beta\left(H+\frac{K}{2}\right)\right)+Q_1^{SC}\right]\right\}.
\end{aligned}
\tag{4.13}
$$

The all possible two-spin correlation functions (beside the $\sigma\sigma$, which is uniform for all HIC and is given by Eq. (3.13)) of the system are listed below,

$$
\begin{aligned}
\langle S^z_{j,1}S^z_{j+r,1}\rangle-\langle S^z_{j,1}\rangle^2 &= \frac{\left(\sum_{k=1}^4\left[A_k^1\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^1\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=1}^4\left[A_k^3\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^3\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}}, \quad (4.14)\\
\langle S^z_{j,2}S^z_{j+r,2}\rangle-\langle S^z_{j,2}\rangle^2 &= \frac{\left(\sum_{k=1}^4\left[A_k^2\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^2\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=1}^4\left[A_k^4\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^4\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}},\\
\langle S^z_{j,1}S^z_{j+r,2}\rangle-\langle S^z_{j,1}\rangle\langle S^z_{j,2}\rangle &= \frac{\left(\sum_{k=1}^4\left[A_k^1\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^1\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=1}^4\left[A_k^4\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^4\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}},\\
\langle S^z_{j,1}\sigma_{j+r}\rangle-\langle S^z_{j,1}\rangle\langle\sigma_j\rangle &= -\frac{e^{-\beta\frac{J}{4}}\sqrt{1-M_\sigma^2}\sum_{k=1}^4\left(A_k^1\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^1\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right)}{4\lambda_1^{SC}}e^{-\frac{r}{\xi}},\\
\langle S^z_{j,2}\sigma_{j+r}\rangle-\langle S^z_{j,2}\rangle\langle\sigma_j\rangle &= -\frac{e^{-\beta\frac{J}{4}}\sqrt{1-M_\sigma^2}\sum_{k=1}^4\left(A_k^2\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^2\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right)}{4\lambda_1^{SC}}e^{-\frac{r}{\xi}},\\
\langle S^z_{j,1}\tau_{j+r}\rangle-\langle S^z_{j,1}\rangle\langle\tau_j\rangle &= \frac{\left(\sum_{k=1}^4\left[A_k^1\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^1\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=0}^4\left[A_k^6\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^6\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}},\\
\langle S^z_{j,2}\tau_{j+r}\rangle-\langle S^z_{j,2}\rangle\langle\tau_j\rangle &= \frac{\left(\sum_{k=1}^4\left[A_k^2\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^2\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=0}^4\left[A_k^6\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^6\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}},\\
\langle\sigma_j\tau_{j+r}\rangle-\langle\sigma_j\rangle\langle\tau_j\rangle &= -\frac{e^{-\beta\frac{J}{4}}\sqrt{1-M_\sigma^2}\sum_{k=0}^4\left[A_k^6\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^6\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]}{4\lambda_1^{SC}}e^{-\frac{r}{\xi}},\\
\langle\tau_j\tau_{j+r}\rangle-\langle\tau_j\rangle^2 &= \frac{\left(\sum_{k=0}^4\left[A_k^5\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^5\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)\left(\sum_{k=0}^4\left[A_k^6\,\mathrm{ch}\left(\beta\frac{kH}{2}\right)+B_k^6\,\mathrm{sh}\left(\beta\frac{kH}{2}\right)\right]\right)}{16\sum_{k=0}^2 D_k\,\mathrm{ch}\left(\beta kH\right)}e^{-\frac{r}{\xi}},
\end{aligned}
$$

where the largest eigenvalue of the transfer-matrix (4.9) is

$$\lambda_1^{SC} = e^{-\beta\frac{J}{4}}\left(\sum_{k=0}^{2} S_k \operatorname{ch}(\beta k H) + \sqrt{\left(\sum_{k=0}^{2} S_k \operatorname{ch}(\beta k H)\right)^2 - 4\sum_{k=0}^{2} D_k \operatorname{ch}(\beta k H)}\right). \tag{4.15}$$

The coefficients in Eqs.(4.14) and (4.15) are

$$\begin{aligned}
&A_1^1 = e^{\beta\frac{3K}{4}} - e^{-\beta\frac{K}{4}}\left(1+2WM_\sigma\right),\ \ A_2^1 = 2e^{\beta\frac{K}{4}}\operatorname{sh}\left(\beta\frac{K}{2}\right)\sqrt{1-M_\sigma^2},\ \ A_3^1 = 0,\ \ A_4^1 = -e^{-\beta\frac{5K}{4}}\sqrt{1-M_\sigma^2}, \quad (4.16)\\
&B_1^1 = \left(e^{\beta\frac{3K}{4}} + e^{-\beta\frac{K}{4}}\right)M_\sigma - 2e^{-\beta\frac{K}{4}}W,\ \ B_2^1 = 2We^{-\beta\frac{K}{4}}\sqrt{1-M_\sigma^2},\ \ B_3^1 = 2e^{-\beta\frac{K}{4}}M_\sigma,\ \ B_4^1 = 0,\\
&A_1^2 = e^{\beta\frac{3K}{4}} - e^{-\beta\frac{K}{4}}\left(1-2WM_\sigma\right),\ \ A_2^2 = A_2^1,\ \ A_3^2 = 0,\ \ A_4^2 = A_4^1,\\
&B_1^2 = \left(e^{\beta\frac{3K}{4}} + e^{-\beta\frac{K}{4}}\right)M_\sigma + 2e^{-\beta\frac{K}{4}}W,\ \ B_2^2 = -2We^{-\beta\frac{K}{4}}\sqrt{1-M_\sigma^2},\ \ B_3^2 = B_3^1,\ \ B_4^2 = 0,\\
&A_1^3 = -e^{\beta\frac{3K}{4}} + e^{-\beta\frac{K}{4}}\left(1-2WM_\sigma\right),\ \ A_2^3 = A_2^1,\ \ A_3^3 = 0,\ \ A_4^3 = A_4^1,\\
&B_1^3 = B_1^2,\ \ B_2^3 = B_2^1,\ \ B_3^3 = B_2^1,\ \ B_4^3 = 0,\\
&A_1^4 = -e^{\beta\frac{3K}{4}} + e^{-\beta\frac{K}{4}}\left(1+2WM_\sigma\right),\ \ A_2^4 = A_2^1,\ \ A_3^4 = 0,\ \ A_4^4 = A_4^1\\
&B_1^4 = B-1^1,\ \ B_2^4 = -B_1^2,\ \ B_3^4 = B_3^1,\ \ B_4^4 = 0,\\
&A_1^5 = \left(2e^{-\beta\frac{K}{4}}Q_2^{SC} + 2e^{\beta\frac{K}{4}}Q_1^{SC} - e^{-\beta\frac{K}{4}} - e^{\beta\frac{3K}{4}}\right)M_\sigma,\ \ A_2^5 = 0,\ \ A_3^5 = 2e^{-\beta\frac{K}{4}}M_\sigma,\ \ A_4^5 = 0\\
&B_0^5 = \left(2e^{\beta\frac{K}{4}}Q_1^{SC} - e^{\beta\frac{3K}{4}}\right)\sqrt{1-M_\sigma^2},\ \ B_1^5 = e^{-\beta\frac{K}{4}}\left(1+2Q_2^{SC}\right) - 2e^{\beta\frac{K}{4}} - e^{\beta\frac{3K}{4}},\\
&B_2^5 = 2\left(e^{\beta\frac{K}{4}}\operatorname{ch}\left(\beta\frac{K}{2}\right) - e^{-\beta\frac{K}{4}}Q_2^{SC}\right)\sqrt{1-M_\sigma^2},\ \ B_3^5 = 0,\ \ B_4^5 = -e^{-\beta\frac{5K}{4}}\sqrt{1-M_\sigma^2}\\
&A_1^6 = A_1^5,\ \ A_2^6 = 0,\ \ A_3^6 = A_3^5,\ \ A_4^6 = 0,\\
&B_0^6 = B_0^5,\ \ B_1^6 = -B_1^5,\ \ B_2^6 = B_2^5,\ \ B_3^6 = 0, B_4^6 = B_4^5,\\
&D_0 = e^{-\beta\frac{K}{2}}\operatorname{sh}^2(\beta K) + \left(e^{\beta K}-1\right)Q_1^{SC} - 2Q_2^{SC}\operatorname{sh}\left(\beta\frac{K}{2}\right),\\
&D_1 = 2e^{-\beta\frac{K}{2}}\left(\operatorname{ch}(\beta K)-1\right)Q_2^{SC} + \operatorname{ch}\left(\beta\frac{3K}{2}\right) - \operatorname{ch}\left(\beta\frac{K}{2}\right),\ \ D_2 = 2Q_2^{SC}\operatorname{sh}\left(\beta\frac{K}{2}\right) + \left(e^{-\beta K}-1\right)Q_1^{SC},\\
&S_0 = e^{\beta\frac{3K}{4}} + 2e^{\beta\frac{K}{4}}Q_1^{SC},\ \ S_1 = 2\left(e^{-\beta\frac{K}{4}}Q_2^{SC} + e^{\beta\frac{K}{4}}\operatorname{ch}\left(\beta\frac{K}{2}\right)\right),\ \ S_2 = e^{-\beta\frac{5K}{4}}.
\end{aligned}$$

V. CONCLUSION

In the present paper we elaborated a technique for calculating correlation functions in HIC. The technique is based on the classical transfer-matric formalism which proves itself as a most efficient and straightforward method of treating the HIC, allowing one to obtain exact expressions for the partition function and, thus, for all thermodynamic functions in a quite simple way[5,9–11,13–17,19–22,26]. However, the classical transfer-matrix formalism is also very convenient for the calculation of the correlation functions of various kind, because all statistical-mechanical expectation values are expressed in terms of traces of certain products of the transfer-matrices with matrices representing the microscopic variables in the transfer-matrix formalism[27]. We presented a detailed scheme of calculation of the correlation function for HIC, obtained general expressions for the corresponding trace formulas and presented the general expressions for the matrix representations of the main microscopic variables for the typical HIC, quantum spin operators, additional classical spins from block Hamiltonians and intermediate spin. One of the main results shows that the spatial decay of any correlation between any pair of spin is characterized by a single correlation length obtained in a standard way as the reciprocal of the logarithm of the ratio of largest and second largest eigenvalues of the classical transfer matrix. In the simplest case considered here when intermediate classical σ-spins are 1/2 spins the transfer matrix has dimension 2 by 2 and, thus, only two eigenvalues. However, the prefactors are strongly affected by the microscopic variables. We illustrated our general formalism on two examples, the simplest HIC, symmetric Heisenberg-Ising diamond-chain and asymmetric Heisenberg-Ising sawtooth chain with a pair of quantum spins in the unit cell. We presented the exact form of the transfer-matrix, its eigenvalues, matrix representation corresponding to the classical transfer-matrix formalism for quantum spin operators, matrix representation for additional classical spin from block Hamiltonian and all possible pair correlation functions. A very important feature of the correlations function of the HIC considered in the present paper is the absence of correlations between transverse components of quantum spins from different blocks. The corresponding matrix representation of the $S^x_{n,1}$ and $S^y_{n,a}$ operators are identically zero. This fact could be connected with the rotational symmetry of the block Hamiltonian with respect to the magnetic field axis.

In is also worth mentioning, that the technique elaborated and illustrated in the present paper, beside of purely theoretical interest, can be used in the practical calculation of the neutron diffraction spectra for the recently obtained coordination-polymer compounds which are the physical examples of HIC[32–34,36,37].

VI. ACKNOWLEDGEMENTS

V.O. expresses his gratitude to the LNF-INFN in Frascati for warm hospitality during the work on the paper. He also acknowledges partial financial support from Volkswagen Foundation (Grant No. I/84 496) and from the project SCS-BFBR 11RB-001.

---

[1] J. Strečka and M. Jaščur, J. Phys.: Condens. Matter **15**, 4519 (2003).
[2] J. Strečka, M. Hagiwara, M. Jaščur, and K. Minami, Czech. J. Phys. **54**, D583 (2004).
[3] J. Strečka, M. Jaščur, M. Hagiwara, K. Minami, Y. Narumi, and K. Kindo, Phys. Rev. B **72**, 024459 (2005).
[4] L. Čanová, J. Strečka and M. Jaščur, J. Phys. : Condens. Matter **18**, 4967 (2006).
[5] J. S. Valverde, O. Rojas, and S. M. de Souza, J. Phys.: Condens. Matter **20**, 345208 (2008).
[6] M. S. S. Pereira, F. A. B. F. de Moura, and M. L. Lyra, Phys. Rev. B **77**, 024402 (2008).
[7] M. S. S. Pereira, F. A. B. F. de Moura, and M. L. Lyra, Phys. Rev. B **79**, 054427 (2009).
[8] J. Strečka, L. Čanová, T. Lučivjanský, and M. Jaščur, J. Phys.: Conf. Ser. **145**, 012058 (2009).
[9] D. Antonosyan, S. Bellucci, and V. Ohanyan, Phys. Rev. B **79**, 014432 (2009).
[10] V. Ohanyan, Condensed Matter Physics **12**, 343 (2009).
[11] V. Ohanyan, Phys. At. Nucl. **73**, 494 (2010).
[12] J. Strečka, L. Galisova, and O. Derzhko, Acta Phys. Pol. A **118**, 742 (2010).
[13] S. Bellucci and V. Ohanyan, Eur. Phys. J. B **75**, 531 (2010).
[14] O. Rojas, S. M. de Souza, V. Ohanyan, and M. Khurshudyan, Phys. Rev. B **83**, 094430 (2011).
[15] O. Rojas and S. M. de Souza, Phys. Lett. A 375, 1295 (2011).
[16] B. Lisnii, Low Temp. Phys. **37**, 296 (2011).
[17] B. Lisnii, Ukr. Journal of Phys. **56**, 1237 (2011).
[18] J. Strečka, M.Dančo, Physica B **406**, 2967 (2011).
[19] V. Ohanyan and A. Honecker, Phys. Rev. B **86**, 054412 (2012).
[20] L. Chakhmakhchyan, N. Ananikian, L. Ananikyan, and Č. Burdık, J. Phys.: Conf. Ser. **343**, 012022 (2012).
[21] O. Rojas, S. M. de Souza, and N. S. Ananikian, Phys. Rev. E **85**, 061123 (2012).
[22] N. Ananikian, L. Ananikyan, L. Chakhmakhchyan, and O. Rojas, J. Phys.: Condens. Matter **24**, 256001 (2012).
[23] T. Verkholyak, J. Strečka, J.Phys. A: Mathematical and Theoretical, **45**, 305001 (2012).
[24] J. Strečka, M. Hagiwara, Y. Han, T. Kida, Z. Honda, and M. Ikeda, arXiv:1206.6656 (unpublished).
[25] L. Galisova, arXiv:1208.0439 (unpublished).
[26] O. Rojas, M. Rojas, N. S. Ananikian and S. M. de Souza, Phys. Rev. A **86**, 042330 (2012).
[27] R. Baxter, *Exactly Solved Models in Statistical Mechanics*, (Academic Press, New York, 1982).
[28] V. Ohanyan and N. Ananikian, Phys. Lett. A **307**, 76 (2003).
[29] E. Aydiner and C. Akyüz, Chin. Phys. Lett. **22**, 2382 (2005).
[30] E. Aydiner, C. Akyüz, M. Gönülol, and H. Polat, phys. stat. sol(b) **243**, 2901 (2006).
[31] D. Gatteschi, R. Sessoli, and J. Villain, *Molecular Nanomagnets* (Oxford Univ. PRess, New York 2007).
[32] D. Visinescu, A. M. Madalan, M. Andruh, C. Duhayon, J.-P. Sutter, L. Ungur, W. Van den Heuvel, and L. F. Chibotaru, Chem. Eur. J. **15**, 11808 (2009).
[33] J. Rinck, G. Novitchi, W. Van den Heuvel, L. Ungur, Y. Lan, W. Wernsdorfer, C. E. Anson, L. F. Chibotaru, and A. K. Powell, Angew. Chem., Int. Ed. **49**, 7583 (2010).
[34] W. Van den Heuvel and L. F. Chibotaru, Phys. Rev. B **82**, 174436 (2010).
[35] V. Ravi Chandra, S. Ramasesha, and D. Sen, Phys. Rev. B **70**, 144404 (2004).
[36] T. S. Venkatakrishnan, S. Sahoo, Nicolas Bréfuel, C. Duhayon, C. Paulsen, A.-L. Barra, S. Ramasesha, and J.-P. Sutter, J. Am. Chem. Soc. **132**, 6047 (2010).
[37] S. Sahoo, J.-P. Sutter, and S. Ramasesha, J. Stat. Phys. **147**, 181 (2012).